

:WhatIsNumerical

2.5 Euler's method

Suppose we have a differential equation of the form

$$\frac{dy}{dt} = f(t, y).$$

We can compare the values of a solution y at times t and $t + \Delta t$ by integrating – the Fundamental Theorem of Calculus tells us that

$$y(t + \Delta t) = y(t) + \int_t^{t+\Delta t} f(\tau, y(\tau)) d\tau. \quad (2.5.1)$$

euler:FTC

Suppose now that we know the value $y(t)$, but do not know the value of the function y for times larger than t . If Δt is not too large, then we can approximate the integral in (2.5.1) using the left endpoint approximation

$$\int_t^{t+\Delta t} f(\tau, y(\tau)) d\tau \approx f(t, y(t)) \Delta t; \quad (2.5.2)$$

euler:left-endpoint-approx

see Figure 2.5.1.




Figure 2.5.1: If Δt is small then the approximation (2.5.2) is “reasonably good.”

fig:left-endpoint-integrat

Using (2.5.2) and (2.5.1) we obtain the approximation

$$y(t + \Delta t) \approx y(t) + f(t, y(t)) \Delta t, \quad (2.5.3)$$

euler:main-approximation

which is “reasonably good” if Δt is small.

We now show how to iteratively use the approximation (2.5.3) in order to construct a numerical approximation of the solution to the initial value problem

$$\frac{dy}{dt} = f(t, y) \quad y(t_0) = y_0. \quad (2.5.4)$$

euler:ivp

We begin by choosing a fixed number Δt and constructing a list of times t_k , indexed by counter k , that each differ by Δt :

k	t_k
0	t_0
1	$t_1 = t_0 + \Delta t$
2	$t_2 = t_0 + 2(\Delta t)$
3	$t_3 = t_0 + 3(\Delta t)$
\vdots	\vdots
k	$t_k = t_0 + k(\Delta t)$

Notice that for each k we have $t_{k+1} = t_k + \Delta t$.

We now construct a list of values y_k that approximate the values of the solution $y(t)$ to (2.5.4) at the times t_k . The initial condition in (2.5.4) provides the value y_0 . The approximation (2.5.3) states that

$$y(t_1) \approx y_0 + f(t_0, y_0)\Delta t.$$

Thus we define

$$y_1 = y_0 + f(t_0, y_0)\Delta t.$$

Similarly, (2.5.3) implies that

$$y(t_2) \approx y(t_1) + f(t_1, y(t_1))\Delta t.$$

We don't know what $y(t_1)$ is, but we do know that $y(t_1) \approx y_1$. Using this approximation we have

$$y(t_2) \approx y_1 + f(t_1, y_1)\Delta t.$$

Thus we set

$$y_2 = y_1 + f(t_1, y_1)\Delta t.$$

Repeating this procedure, we see that the numbers y_k , defined by

$$y_k = y_{k-1} + f(t_{k-1}, y_{k-1})\Delta t, \quad (2.5.5)$$

euler:euler

approximate the value of $y(t_k)$. The approximation scheme (2.5.5) is called ***Euler's method***.

We can summarize this in the following table:

k	t_k	y_k
0	t_0	y_0
1	$t_1 = t_0 + \Delta t$	$y_1 = y_0 + f(t_0, y_0)\Delta t$
2	$t_2 = t_0 + 2(\Delta t)$	$y_2 = y_1 + f(t_1, y_1)\Delta t$
3	$t_3 = t_0 + 3(\Delta t)$	$y_3 = y_2 + f(t_2, y_2)\Delta t$
\vdots	\vdots	
k	$t_k = t_0 + k(\Delta t)$	$y_k = y_{k-1} + f(t_{k-1}, y_{k-1})\Delta t$

Let's now look at how this approximation works for an equation for which we know the exact solution.

Example 2.5.1. Consider the initial value problem

$$\frac{dy}{dt} = y^2 \quad y(0) = 1.$$

We know that the solution to the initial value problem is

$$y(t) = \frac{1}{1-t}. \quad (2.5.6)$$

Let's now construct an approximation using Euler's method.

We choose $\Delta t = 0.1$ and decide to approximate the solution $y(t)$ on the interval $0 \leq t \leq 0.5$. Since $t_0 = 0$, we are interested in the times

k	t_k
0	$t_0 = 0$
1	$t_1 = 0.1$
2	$t_2 = 0.2$
3	$t_3 = 0.3$
4	$t_4 = 0.4$
5	$t_5 = 0.5$

We now use (2.5.5) in order to construct the list of approximate values. Since $f(t, y) = y^2$ and $y_0 = 1$, we compute

$$y_1 = y_0 + f(t_0, y_0)\Delta t = 1 + (0.1)^2(0.1) = 1.001.$$

Proceeding iteratively, we obtain the following table:

k	t_k	y_k
0	$t_0 = 0$	$y_0 = 1$
1	$t_1 = 0.1$	$y_1 = 1.100$
2	$t_2 = 0.2$	$y_2 = 1.221$
3	$t_3 = 0.3$	$y_3 = 1.370$
4	$t_4 = 0.4$	$y_4 = 1.558$
5	$t_5 = 0.5$	$y_5 = 1.800$

A plot of the points (t_k, y_k) , along with a plot of the solution, appears in Figure 2.5.2.




Figure 2.5.2: A plot of the approximating points (t_k, y_k) , as computed with $\Delta t = 0.1$, in red and a plot of the actual solution $y(t)$ in blue. Notice that the error accumulates as time progresses.

fig:euler-exampl

As evidenced by the plot in Figure 2.5.2, the approximation scheme with $\Delta t = 0.1$ is not bad, but is not great. We can obtain a better approximation by choosing $\Delta t = 0.01$, in which case we must plot 50 points. The comparison between this better approximation and the actual solution appears in Figure 2.5.3.

We can easily make Sage do the computations in Euler's method. Here I explain how to build the code used in Example 2.5.1. First we define variable y and also the function $f(y) = y^2$ that determines the right hand side:

```
var('y')
f(y) = y^2
```

Then we define Δt and set it equal to 0.1. We also define y_0 , which we set equal to 1:

```
var('y')
f(y) = y^2
```




Figure 2.5.3: A plot of the approximating points (t_k, y_k) in red, now with $\Delta t = 0.01$, together with a plot of the actual solution $y(t)$ in blue. Notice that the approximation here is much better than the approximation in Figure 2.5.2.

fig:euler-example2

```
y0 = 1
deltaT = 0.1
```

Next we construct a data structure that will hold the various values of t_k and y_k . To do this we make a variable called `steps` that tells us how many time steps we will take. In this example, we set `steps` equal to 5. We call that data structure `eulerData`. For now, we define all the t_k values to be $t_k = k(\Delta t)$ and all the y_k values to be the same as y_0 .

```
var('y')
f(y) = y^2

y0 = 1
deltaT = 0.1
steps=5

eulerData = [[k*deltaT,y0] for k in range(0,steps+1)]

eulerData
```

The variable `eulerData` is organized as follows: the quantity `eulerData[2]` tells us the entries in the row corresponding to $k = 2$.

```

var('y')
f(y) = y^2

y0 = 1
deltaT = 0.1
steps=5

eulerData = [[k*deltaT,y0] for k in range(0,steps+1)]

eulerData[2]

```

If we want only to see the time t_2 , then we need to ask Sage to show us `eulerData[2][0]`, while if we want Sage to show us y_2 , we need to ask to be shown `eulerData[2][1]`. Try it:

```

var('y')
f(y) = y^2

y0 = 1
deltaT = 0.1
steps=5

eulerData = [[k*deltaT,y0] for k in range(0,steps+1)]

eulerData[2][0]

```

What we want to do now is systematically go through and update all the y_k values, starting at $k = 1$ and ending at $k = \text{steps}$. We can do this with a loop:

```

var('y')
f(y) = y^2

y0 = 1
deltaT = 0.1
steps=5

eulerData = [[k*deltaT,y0] for k in range(0,steps+1)]

for k in [1..steps]:
    eulerData[k][1]=eulerData[k-1][1]+deltaT*f(eulerData[k-1][1])

eulerData

```

Finally, we can plot the resulting approximate solution:

```

var('y')

```

```

f(y) = y^2

y0 = 1
deltaT = 0.1
steps=5

eulerData = [[k*deltaT,y0] for k in range(0,steps+1)]

for k in [1..steps]:
    eulerData[k][1]= eulerData[k-1][1] + deltaT*f(eulerData[
        k-1][1])

eulerPlot = list_plot(eulerData, color="red", plotjoined=
    true,marker='.',axes_labels=['t$', '$y$'])
eulerPlot.show()

```

Activity 2.5.1. Use Euler's method to construct an approximate solution to the initial value problem

$$\frac{dy}{dt} = 2 \cos(y), \quad y(0) = \frac{1}{2}.$$

Plot the numerical solution on top of the slope field.

TableForTwo

Exercise 2.5.1. Create an approximate table of values for the solution of the IVP

$$\frac{dy}{dt} = \cos(y), \quad y(0) = 0$$

over the interval $0 \leq t \leq 1$. Please focus on the step-size of $\Delta t = 0.25$.

WNM-MidEnd

Exercise 2.5.2. As you know, the function $y(t) = e^t$ is the unique solution of the IVP

$$\frac{dy}{dt} = y, \quad y(0) = 1.$$

In this problem you get to investigate the power and the limitation of the Euler method.

1. Make a table which shows the actual values of $y(0)$, $y(0.25)$, $y(0.5)$, $y(0.75)$ and $y(1)$.
2. What does the Euler method for step-size of $\Delta t = 0.25$ predict for the values of $y(0)$, $y(0.25)$, $y(0.5)$, $y(0.75)$ and $y(1)$?
3. What does the Euler method for step-size of $\Delta t = 0.05$ predict for the values of $y(0)$, $y(0.25)$, $y(0.5)$, $y(0.75)$ and $y(1)$?

4. Make a table that lists the errors that the two approximation methods make. Then write a sentence or so interpreting what you see.
5. Represent your findings graphically in the t - y plane. Then write a sentence or so interpreting what you see.

WMM-MidBegin

Exercise 2.5.3. Consider the differential equation

$$\frac{dy}{dt} = y(y - 3)(y - 6).$$

1. Perform qualitative analysis of this equation.
2. In addition to the differential equation above consider the initial condition $y(0) = 1$. Perform Euler method with the step size $\Delta t = 0.5$ in order to understand $y(t)$ for $0 \leq t \leq 2$. Make sure your answer includes the table of values and the corresponding piece-wise linear graph.
3. In light of part (a) please discuss the reliability of your answer in part (b).

HowAreYouTodayMyFriend

Exercise 2.5.4. Consider the initial value problem

$$\frac{dy}{dt} = y^3, \quad y(0) = 1.$$

1. Using the Euler's Method with $\Delta t = 0.5$, graph an approximate solution over the interval $0 \leq t \leq 1$.
2. What happens if you make Δt considerably smaller than above (e.g. $\Delta t = 0.05$)?
3. Verify that the function

$$y(t) = \frac{1}{\sqrt{1 - 2t}}$$

is a solution to the IVP in this problem. Use this knowledge to interpret the findings in parts a) and b) of this problem.

WMM-Extra-Only

Exercise 2.5.5. Construct a numerical solution to the initial value problem

$$\frac{dy}{dt} = -y^2 + t, \quad y(2) = 2.$$

Use your solution to estimate $y(10)$.